

Biases in Expansion Distances of Novae Arising from the Prolate Geometry of Nova Shells

Richard A. Wade^{1,2}

Jason J. B. Harlow^{1,3}

and

Robin Ciardullo^{1,4}

ABSTRACT

Expansion distances (or expansion parallaxes) for classical novae are based on comparing a measurement of the shell expansion velocity, multiplied by the time since outburst, with some measure of the angular size of the shell. We review and formalize this method in the case of prolate spheroidal shells. For such shells there is no unique angular size except when the shell is seen pole-on, and several different measures of angular size have been used in the literature. We present expressions for the maximum line-of-sight velocity from a complete, expanding shell and for its projected major and minor axes, in terms of the intrinsic axis ratio and the inclination of the polar axis to the line of sight. For six distinct definitions of “angular size”, we tabulate the error in distance that is introduced under the assumption of spherical symmetry (i.e., without correcting for inclination and axis ratio). The errors can be significant and systematic, and can affect studies of novae whether considered individually or statistically. Each of the six estimators overpredicts the distance when the polar axis is close to the line of sight, and most underpredict the distance when

¹Department of Astronomy & Astrophysics, The Pennsylvania State University, 525 Davey Laboratory, University Park, PA 16802-6305

²wade@astro.psu.edu

³Present address: Department of Physics, University of the Pacific, 3601 Pacific Ave., Stockton, CA 95211-0197. email: jharlow@uop.edu

⁴rbc@astro.psu.edu

the polar axis is close to the plane of the sky. Use of the straight mean of the projected semimajor and semiminor axes gives the least distance bias for an ensemble of randomly oriented prolate shells, and we recommend this method when individual inclinations and axis ratios cannot be ascertained. The best individual expansion distances, however, result from a full spatio-kinematic modeling of the nova shell. We discuss several practical complications that affect expansion distance measurements of real nova shells. We recommend that nova shell expansion distances be based on velocity and angular size measurements made contemporaneously if possible, and that the same ions and transitions be used for the imaging and velocity measurements. We emphasize the need for complete and explicit reporting of measurement procedures and results, regardless of the specific method used.

Subject headings: novae — circumstellar matter

1. INTRODUCTION

The distance to a classical nova in the Galaxy is best inferred by comparing the angular size of the resolved nova shell with the size calculated from its rate of expansion and the time since the shell was ejected. However, if nova shells are ellipsoids of revolution (spheroids) rather than spherical, then the concept of “angular size” is ambiguous, and the expansion velocity along the line of sight does not correspond to the transverse expansion velocity. Thus the use of formulas that are valid in the spherical case will lead to erroneous distance estimates. Individual distance estimates may be too large or too small, depending on the true axis ratio of the nova shell and its inclination to the line of sight. Furthermore, these errors do not necessarily average toward zero when ensemble averages are taken. In this paper, we consider systematic errors in estimates of nova shell expansion distances, and recommend procedures to minimize the errors.

1.1. The Usefulness of Nova Distances

As with all classes of astronomical objects, our understanding of the classical nova phenomenon depends on having accurate estimates of the distances of these objects. In turn, having a well-founded understanding of the distances and luminosities of novae allows them to be studied as astrophysical objects and to be exploited for other purposes.

The need for accurate distances is evident, both for novae taken individually and for novae used collectively, i.e., in a statistical fashion. An accurate distance to an individual nova, combined with good coverage of the outburst light curve and a knowledge of the interstellar extinction, can allow the theory of the nova outburst to be verified and further developed. For example, it is possible to check whether there is a phase after maximum light during which the nova’s luminosity is close to the Eddington limit. All inferences about the mass of the shell that is ejected during the outburst depend on some power of the distance, through establishing the volume occupied by the emitted gas. At late stages in the evolution of a nova, when the shell can be resolved from the central binary star, the distance is needed to convert the angular size of the shell into a linear size, so that the physical conditions in the ejected gas (ionization, excitation) can be related to the ionizing flux from the central white dwarf (the post-nova) and its accretion disk. An accurate distance to a classical nova also allows the modeling of the accretion process in the post-nova system; without a distance constraint (leading to a constraint on the luminosity), it has proven impossible to infer uniquely the mass accretion rate onto the white dwarf (Wade 1988, Wade & Hubeny 1998).

Treated collectively, novae have the possibility of providing a secondary or even primary distance indicator for the extragalactic distance scale. Since novae are present in galaxies of all Hubble types, they have the potential to be used directly to compare and unite the distance scales of spiral and elliptical galaxies. The so-called maximum magnitude – rate of decline (MMRD) relation gives the visual absolute magnitude at maximum light, from a measurement of the rate of decline after maximum (or equivalently the time taken to decline 2 or 3 magnitudes). The shape of the mean MMRD curve and the dispersion around this mean relation have been found from observations in external galaxies (e.g., Della Valle & Livio 1995). For the MMRD relation to be a primary distance indicator, however, the zero-point calibration must be provided by Galactic novae. Another proposed distance indicator is the absolute magnitude at 15 days past maximum light (M_{15}), where the dispersion in absolute magnitude is small for all novae taken without regard to speed class. The same remarks about zero-point calibration apply to this method.⁵

Even the nearest Galactic novae are generally too distant for direct trigonometric parallax measurements. Instead, indirect methods of distance estimation are often used, based on the Galactic rotation curve, the total amount of interstellar reddening, the presence or absence of discrete components (“clouds”) in interstellar absorption lines, etc.

⁵As one step in calibrating the MMRD relation, Cohen (1985) adjusted M_V at maximum for her best observed novae, assuming that they had *identical* M_{15} ; this step should be replaced by actually measuring the dispersion in M_{15} .

The only geometrical (hence fundamental) method is that of “expansion distance” (also referred to as “expansion parallax”), in which the measured angular size of the resolved nova shell is compared with the linear size of the shell; the latter is calculated from the expansion speed of the shell gas and the known time since the outburst. In her work on the MMRD relation, Cohen (1985) had only eleven novae with well-observed expansion distances and suitable coverage of their light curves. This is largely because the surface brightness of a nova shell declines rapidly with time since outburst, so that by the time the shell is large enough to be resolved from the ground, it is often too faint to observe. Since Cohen’s study, several additional novae have had good light curve coverage, and expansion distances may become available for these from ground-based observations and especially from the *Hubble Space Telescope*⁶, since the latter can resolve some shells within a few months or years of the outburst.

The simplest way to derive an expansion distance is to assume (often implicitly) that the nova shell is expanding spherically symmetrically, hence that the transverse velocity of gas in the plane of the sky is the same as the radial velocity of gas moving directly along the line of sight. What if the ejection of the shell is asymmetric? To be specific, suppose the shell expands as a spheroid, the simplest generalization from the spherically symmetric case, and one that suffices to describe many actual nova shells. First, the projected image of the nova shell will not be circular for most orientations, and thus there will be an ambiguity in what is meant by angular size. Second, the maximum velocity along the line of sight will usually not correspond to *either* the “polar” or “equatorial” expansion velocity. For example, suppose that the angular size is taken to be the largest projected “radius” of the nova shell, which is perhaps the easiest size parameter to estimate on a barely resolved image. If all nova shells were oblate, then the calculated expansion distance (based on the assumption of spherical symmetry) would always be less than the true distance, because the line-of-sight velocity would be smaller than the transverse expansion velocity. The resulting nova distance scale would be too short. On the other hand, if all nova shells are prolate, then the distance to an individual object may be underestimated or overestimated, depending on the orientation and the ratio of major and minor axes. While it is clear that the distance to an individual nova can be in error as a result, it was not made clear until the work of Ford & Ciardullo (1988; hereinafter FC88) that in the prolate case, a systematic error might remain, even after averaging over an ensemble of novae that are taken to be randomly oriented in space. In their analysis, FC88 made the assumption stated above as

⁶Narrow-band imagery of recent novae has been carried out with the HST Wide Field/Planetary Camera 2, used in “snapshot” mode, as part of program 7386; these images are public. HST imagery of somewhat older nova shells is discussed in Gill & O’Brien (2000).

an example, that the angular size of the nova shell is taken to be the major axis of the projected image. However, all workers do not make this identification. For example, Cohen & Rosenthal (1983) did use the projected semimajor axis for the angular size, but Cohen (1985) used an angle-averaged radius. What way is best? A goal of this paper is to extend the FC88 analysis to include consideration of six distinct yet plausible ways of defining the angular size of the shell.

As the number of Galactic novae with well observed light curves increases, and with the much greater resolving power provided by adaptive optics and HST, it is likely that the calibration of the MMRD and M_{15} relations will be improved, but the question of possible systematic errors in the distances becomes more important. This is especially so, if shell morphology is related to nova speed class, as has been suggested by Slavin, O’Brien, & Dunlop (1995). Likewise, as more expansion distances for individual novae become available, it is important to have clearly in mind whether and how much these distances may be in error, as the result of measuring uncertainties and modeling assumptions.

1.2. Prolate or Oblate?

Theoretical arguments have been made favoring both oblate and prolate geometries for nova shells (Porter, O’Brien, & Bode 1998 and references therein). Empirically, it is now the consensus that, to the extent that nova shells can be described by spheroids, they are either prolate or spherical, but not oblate. FC88 discussed the few cases of resolved nova shells known at the time, in terms of whether they were elongated along one axis (prolate spheroids) or compressed along one axis (oblate spheroids). FC88 noted that most data on the shapes of nova shells were consistent with spherical or prolate geometries, but categorized the shell of nova HR Del 1967 as oblate. For this object, early models by Hutchings (1972) and Soderblom (1976) indeed suggested an oblate symmetry. A spatio-kinematic model by Solf (1983), however, has clearly shown that the resolved shell of HR Del is consistent with a prolate geometry, and not consistent with being oblate. Slavin, O’Brien, & Dunlop (1995) carried out imaging of nova shells using narrow band filters; in particular they have obtained images at several different tilts of an interference filter with nominal wavelength 6560 Å (17 Å FWHM), which allowed them to distinguish crudely between gas approaching or receding from the observer. Their data are clearly consistent with the shells being prolate, not oblate, if they depart detectably from spherical symmetry. Other spatio-kinematic studies, for example of the shell around nova DQ Her 1934 (e.g. Herbig & Smak 1992) also indicate prolate symmetry. Therefore we proceed with the assumption that to first approximation, nova shells are prolate spheroidal shells, with

their properties in projection specified by their axis ratio and the inclination of the polar axis to the line of sight.

In Section 2 of this paper, we investigate several different ways of defining the angular size of the resolved nova shell. We derive the projected size and shape of the shell and the maximum radial velocity of gas in the shell, as functions of the intrinsic axis ratio and the inclination of the polar axis. We then derive analytic expressions that give the inferred distance in terms of the true distance, as a function of axis ratio and inclination. We tabulate results for a variety of cases, using six definitions of “angular size.” In Section 3 we investigate how these various definitions of expansion distance behave, both for individual objects and when averaged over an ensemble of nova shells oriented randomly in space. We also discuss some practical matters relating to the measurement of the expansion speed and angular size of nova shells. We summarize our findings in Section 4.

2. EXPANSION DISTANCE ESTIMATORS FOR PROLATE SPHEROIDAL NOVA SHELLS

When viewed at an inclination angle i , a prolate spheroid at distance d and with principal axis ratio $b/a \leq 1$ will appear projected on the sky as an ellipse with apparent axis ratio $b_{\perp}/a_{\perp} \leq 1$. We define several quantities, which appear repeatedly in the discussion to follow:

$$\begin{aligned} f_1 &= \sqrt{1 - e^2 \sin^2 i} \\ f_2 &= \sqrt{1 - e^2 \cos^2 i} \\ f_3 &= \sqrt{1 - e^2} = b/a \end{aligned}$$

The auxiliary quantity e is the “eccentricity” of the prolate spheroid, in the sense that $b^2 = a^2(1 - e^2)$ relates the major and minor axes, a and b respectively, of an ellipse.

We have the following relations between the axes of the spheroid in space, a and b , and the (linear) principal axes of the projected ellipse, $a_{\perp}$ and $b_{\perp}$ (see Appendix A).

$$\begin{aligned} a_{\perp} &= f_2 a \\ b_{\perp} &= b = f_3 a = (f_3/f_2) a_{\perp} \end{aligned}$$

Let v_0 denote the expansion speed along the major (polar) axis of the spheroid. Then from Appendix B the maximum projected (line-of-sight) speed is

$$v_{\max} = f_1 v_0 = f_1(a/t),$$

where a is the semimajor axis of the spheroid when the age of the nova remnant is t . (Constant expansion speed is assumed.) Also, a distance x measured in the plane of the sky corresponds to an angle $\rho_x = x/d$ (radians), where d is the true distance of the nova.

The essence of the expansion distance method is to compare an estimate of the linear size of the nova shell, vt , with an estimate of the angular size, ρ . The estimator formula

$$\hat{d} = \frac{v_{\max}t}{\rho}, \quad (1)$$

where ρ is any angular radius, recovers the true distance d in the case of a spherically symmetric expanding shell. This is because $v_{\max} = v_0$ (by symmetry) and all angular radii are equal to a/d where a is the true linear size of the shell at time t :

$$\hat{d} = \frac{v_{\max}t}{\rho} = \frac{v_0t}{(a/d)} = d. \quad (2)$$

For a prolate spheroid, in general $v_{\max} \neq v_0$, and there is no unique measure of the angular size ρ . Given independent knowledge of i , the apparent ratio $b_{\perp}/a_{\perp} = f_3/f_2 = (1 - e^2)^{1/2}/(1 - e^2 \cos^2 i)^{1/2}$ can be inverted to find e . In this case equation (1) can be used, with corrections to convert the measured speed $v_{\max}$ into v_0 and the apparent semimajor axis $\rho_1 = a_{\perp}/d$ into a/d , to recover the true distance. In symbols,

$$\hat{d} = \frac{(v_{\max}/f_1)t}{(\rho_1/f_2)} = \frac{v_0t}{(a/d)} = d. \quad (3)$$

In general, however, the inclination of the spheroid in space is not known, so the correction factors are not known, but the simple formula $\hat{d} = v_{\max}t/\rho_1$ does not recover d . Furthermore, there is no reason any longer to define ρ as the apparent semimajor axis — the apparent minor axis or some average indicator of the apparent size of the nova shell could be used instead.

We consider six possible definitions of ρ , and for each we investigate how large an error is made in estimating d using equation (1). This question is addressed both for individual novae, with particular values of b/a and i , and for statistical ensembles of nova shells, where averages are taken over random orientations for a fixed value of b/a .

The six choices for ρ are:

$$\begin{aligned}
\rho_1 &= a_{\perp}/d &= f_2 \times a/d \\
\rho_2 &= b_{\perp}/d &= f_3 \times a/d \\
\rho_3 &= (a_{\perp} + b_{\perp})/2d &= (f_3 + f_2)/2 \times a/d \\
\rho_4 &= \sqrt{a_{\perp} b_{\perp}}/d &= \sqrt{f_3 f_2} \times a/d \\
\rho_5 &= 2b_{\perp}K(k)/\pi d &= 2f_3K(k)/\pi \times a/d \\
\rho_6 &= 2(a_{\perp}^{-1} + b_{\perp}^{-1})^{-1}/d &= 2(f_2^{-1} + f_3^{-1})^{-1} \times a/d
\end{aligned}$$

The first two choices are the apparent major and minor axes of the projected ellipse. The third choice is the arithmetic mean of ρ_1 and ρ_2 . The fourth choice is the geometric mean of these. The fifth definition, ρ_5 , is the angle-averaged apparent “radius” of the shell. Here $K(k)$ is the complete elliptic integral of the first kind (see Appendix C). The argument k is given by $k^2 = 1 - (k')^2 = 1 - (b_{\perp}/a_{\perp})^2 = 1 - (f_3/f_2)^2$. The sixth choice is the harmonic mean of ρ_1 and ρ_2 .

Corresponding to each ρ_i is a distance estimator $\hat{d}_i = v_{\max}t/\rho_i$. Given in Table 1 are $k' = b_{\perp}/a_{\perp}$, $v_{\max}/v_0$, $a_{\perp}/a$, and $\hat{d}_j/d$, ($j = 1, 2, \dots, 5$) for several combinations of true axis ratio b/a and inclination i . Since $\hat{d}_6/d = (\hat{d}_1 + \hat{d}_2)/2d$, it is not shown separately in Table 1.

The probability that the polar axis of a randomly oriented spheroid makes an angle between i and $i + di$ with the line of sight is $P(i)di = (\sin i)di$. Each of the estimates $\hat{d}_j$ can be averaged over angle with $P(i)$ as the weighting function:

$$\langle \hat{d}_j \rangle = \int_0^{\pi/2} P(i) \hat{d}_j(i) di. \quad (4)$$

The $P(i)$ -weighted average values of $\hat{d}_j/d$ were computed numerically and are shown in Table 1. For example, $\langle \hat{d}_1 \rangle/d = 0.937$ for $b/a = 0.80$.

3. RESULTS AND DISCUSSION

3.1. The Typical Size of Errors in Expansion Distances

Inspection of Table 1 reveals several features of the various distance estimators $\hat{d}_i$. Every estimator gives the correct distance for spherically symmetric nova shells, as expected. For prolate nova shells, every estimator overpredicts the distance when the polar axis is close to the line of sight, because $v_{\max}$ does not correspond to the projected angular size. In this case, all estimators are equally poor, and increasingly so as the axis ratio b/a becomes more extreme. All estimators except $\hat{d}_2$ underpredict the distance for prolate nova shells

when the polar axis is close to the plane of the sky. In this case, the error increases as the axis ratio decreases, but $\hat{d}_5$ and $\hat{d}_6$ are better than the other estimators.

Finally, when considering an ensemble average of shells with random orientations, the best average distance is produced using $\hat{d}_3$, which is based on the straight mean of the projected semimajor and semiminor axes. Since this is true for each intrinsic axis ratio considered separately, it will also be true for an ensemble of randomly oriented nova shells that has a mixture of axis ratios.

Observations of resolved nova shells show a variety of projected axis ratios. For example, nova DQ Her 1934 has a projected axis ratio of $k' = b_{\perp}/a_{\perp} = 0.73$ (Herbig & Smak 1992), while nova FH Ser 1970 has $k' = 0.91$ (Slavin, O’Brien, & Dunlop 1995). Nova HR Del 1967 has $k' = 0.56$ according to Solf (1983). Slavin et al. (1995) find $k' = 0.75$ for HR Del viewed in H α , but a more elongated image (same projected major axis, shorter projected minor axis) in the light of [O III].⁷ Solf finds the intrinsic axis ratio for HR Del to be $b/a = 1/3$, possibly the most extreme of the well-studied shells. Thus Table 1 covers most of the axis ratios observed to date. If there were a shell with $b/a = 0.4$ viewed pole-on, the expansion distance method would yield an estimated distance a factor of 2.5 larger than the true distance, regardless of the specific estimator used. The median inclination of the polar axis is 60° , and 80 per cent of all randomly oriented nova shells should have inclinations between 26° and 84° . Thus *typical* individual errors in nova expansion distances will be of order tens of percent, unless the inclination and axis ratio are known. Barring the tell-tale evidence of high inclination, given by an eclipsing binary system, the only way to derive the inclination of a nova shell is to construct a so-called spatio-kinematic model, in which spectra from multiple positions across the resolved shell are simultaneously modeled (e.g., HR Del, Solf 1983; DQ Her, Herbig & Smak 1992, Gill & O’Brien 2000).

3.2. Practicalities of Measuring Nova Shells

The nova shell literature contains a mixture of techniques for measuring the expansion velocities and angular sizes. Given that nova shells typically are bright enough to detect only for a few decades after outburst and hence have small angular sizes if they are resolved,

⁷Slavin, O’Brien, & Dunlop (1994) discuss the different appearance of HR Del in different ions as perhaps arising from density or composition differences of the gas ejected in different directions from the central star. Another possibility is a difference in ionizing flux between the poles and the equator of the shell, with the equator being shielded by the accretion disk around the star (cf. the discussion of nova FH Ser 1970 by Gill & O’Brien 2000).

this is not surprising. Nevertheless, it is clear that velocities and angular sizes have not always been combined in a self-consistent fashion, even in cases where the inclination and axis ratio are known. Herbig & Smak (1992) discuss this point at length for the case of nova DQ Her 1934, comparing several different spatio-kinematic studies that arrived at widely differing distance estimates. For DQ Her the issue is that the shell has a finite thickness; it is therefore possible erroneously to combine a velocity measured from the extreme outer edge with, say, a size measured from the ridge line of an image, which represents a position *within* the shell. Martin (1989) also discusses this point. To avoid confusion, it is highly desirable for observers reporting either a line-of-sight expansion velocity or an angular size to be explicit about exactly what was measured. We suggest that angular size measurements be referred to a contour level that encloses a stated percentage of the nova shell flux. (Care needs to be taken, if imaging is done through a narrow-band filter that excludes part of the shell emission.) Likewise, emission line velocities may be measured at a level above the continuum such that a stated percentage of the total line flux is at less extreme velocities.

If image deconvolution methods (e.g., the MEM and CLEAN algorithms) are used to “sharpen” the image, special care needs to be taken in determining and reporting any effect this has on measurements of the “edge” of a nova shell, especially if the method does not conserve flux.

In addition to avoiding possible ambiguities caused by the thickness of the shell, authors need to be clear about what sort of angular size is being reported. Cohen & Rosenthal (1983) used ρ_1 , the projected semimajor axis of the shell, and FC88 modeled the systematic errors in nova expansion distances assuming ρ_1 , as it is perhaps the easiest to measure for barely resolved shells. Cohen (1985), however, derived angular sizes by carefully modeling the appearance of a (spherically symmetric) shell superimposed on a central star, taking into account the point spread function of the optical system. This technique gives ρ_5 . Others, e.g., Shin et al. (1998), have been less careful, merely measuring the FWHM of the nova image and subtracting a stellar FWHM in quadrature to obtain a “characteristic” size. While this method is adequate to demonstrate angular extension of the shell, it is essentially useless for distance estimation, since it wrongly assumes that the profile of a seeing-convolved image of a shell is gaussian, and it does not take into account light from the central star.

Additional complications arise if the projected image of the nova shell is elongated but not strictly elliptical, or if the outline of the shell is incomplete. Herbig & Smak (1992) demonstrated that the “equator” of the DQ Her shell is constricted in both angular extent and velocity, and were careful to distinguish the measured equatorial velocity from the modeled minor-axis velocity of the spheroid. If there is emission from gas *beyond* the main

elliptical outline, as reported for DQ Her (Slavin, O’Brien, & Dunlop 1995) and RR Pic (Gill & O’Brien 1998), care must be taken that the extreme velocity used to estimate distance corresponds to the shell from which the angular size was taken, and not the extended halo. Furthermore, if the spheroidal shell is incomplete, consisting only of an equatorial “belt” and polar “blobs”, then for most inclination angles, there will be no emitting gas at the internal angle θ (see Figure 2) that would correspond to v_{max} in a filled shell. Likewise, there may not be any gas emitting at the tangent to the line of sight, which defines the projected major axis for a complete shell (Appendix A). In such a case, our prescription for distance estimators formally breaks down. However, FC88 give a partial discussion of this case in the limit of a narrow equatorial belt and small polar caps, giving expressions for $\hat{d}_1/d$ and its average for random orientations. FC88 find that individual nova distances for the “belt and blobs” case can be over- or underestimated, up to the same extremes as for the complete shell case, and the angle-averaged distance is underestimated by an amount very similar to the complete shell case discussed here.

Another complication arises from the fact that some nova shells have been observed to change their projected shape in the few years immediately after outburst. A notable case is that of nova V1974 Cyg 1992, which was observed with long-baseline interferometry at radio frequencies (Hjellming 1996). The radio data suggest a model in which the outer and inner faces of the expanding shell have different (triaxial) ellipsoidal representations. Moreover, careful models constructed to account for radio observations of novae (Hjellming et al. 1979; Seaquist et al. 1980) demonstrate that the entire expanding gas cloud does not contribute equally to the optical emission line radiation, since the shell is likely more dense at the inner boundary. The emissivities of the various ions and the fraction of the expanding shell that contributes to the flux both change with time, as the nova shell expands and the post-nova central binary (which photoionizes the shell) recovers from the eruption. The size and morphology of a given nova shell image may differ, depending on which transition is used to image it. As discussed in Section 3.1, HR Del differs considerably in appearance, depending on whether the shell is imaged in $\lambda 5007$ [O III] or in $\lambda\lambda 6548/6563/6584$ H α + [N II]. For all these reasons, it is best to measure the expansion velocity and the angular size of a shell contemporaneously if possible; this will assure that the same parcels of gas are used for both measurements.

Finally, since the central star of a nova system is a cataclysmic binary, it is possible to conflate an emission line from an accretion disk or magnetically channeled accretion flow in the binary with the same line (or a nearby one) from gas in the nova shell. Since orbital speeds of gas in the inner accretion disk may be of order 10^3 km s $^{-1}$, there is the possibility to mis-measure the expansion velocity of the shell, unless care is taken to measure v_{max} from a long-slit spectrum oriented along the projected major axis. (Recall from Appendix

B that the point of the projected image at which v_{max} is attained is generally not the same point as the central star.)

These examples of traps and complications in the estimation of nova expansion distances are intended to reinforce a plea that observers give a full description of their methods and results. Depending on the particular nova, the information available, and the goals of the study, one method of proceeding or another may be appropriate — perhaps there is no single way to estimate an expansion distance that is best in every instance. To allow results from different studies to be combined correctly, the need for careful documentation is evident.

4. SUMMARY AND CONCLUSION

We have reviewed and formalized the method of nova shell expansion distances as a means of estimating the distances to classical novae. This method combines a measurement of the shell expansion velocity (multiplied by the time since outburst) with some measure of the angular size. Expansion distances for novae underlie the calibration of the MMRD and M_{15} relations and also form the basis for astrophysical studies of individual novae and their remnants. It is therefore important to adopt methods of measurement that minimize any possible bias in the distances that results from incomplete information about the shape or orientation of the nova shells.

Many resolved shells exhibit significant prolate symmetry, so that there is no unique angular size except when the shell is seen pole-on. We developed analytic expressions for the maximum line-of-sight velocity from a complete, expanding prolate spheroidal shell and for its projected major and minor axes, as functions of the intrinsic axis ratio and the inclination of the polar axis to the line of sight. For six definitions of “angular size”, we then computed the error introduced by deriving a distance using the assumption of spherical symmetry (i.e., without correcting for inclination and axis ratio). The errors can be significant and possibly systematic, affecting studies of novae whether considered individually or statistically.

The definition of angular size that results in the least errors at the extremes is ρ_6 , the harmonic mean of the projected semimajor and semiminor axes. However, the definition that results in the least bias when an ensemble of randomly oriented prolate shells is considered is ρ_3 , the straight mean of the projected semimajor and semiminor axes, and we recommend this method when individual inclinations and axis ratios cannot be ascertained. The ρ_3 -based method is always as good or better than the ρ_1 method

(projected semimajor axis alone). The best individual expansion distances result from a full spatio-kinematic modeling of the nova shell, using spectroscopy of emission lines at multiple locations across the resolved shell.

We have discussed practical issues and made recommendations for observers who make measurements of either the maximum line-of-sight velocity and the angular size of a resolved nova shell. The velocity measurement may be complicated by the presence of line emission from the central cataclysmic binary star, and if the spheroidal shell is not complete, the theoretical maximum velocity may not be observed at all. The correct application of angular size measurements can be compromised by convolution with the image point spread function, by improper technique, or by incomplete reporting. For best results, velocity and angular size measurements should be made contemporaneously, and must refer to the same features of the shell. Observers are encouraged to report as completely as possible the measurements they have made. Estimates of nova distances by the shell expansion method (or any other method) should be accompanied by a discussion of both random and systematic errors, including possible effects due to unaccounted-for departures from spherical symmetry, as discussed in this paper.

Support for this work was provided by NASA through grant number GO-07386.01 from the Space Telescope Science Institute, which is operated by AURA, Inc. under NASA contract NAS 5-26555. This research has made use of the Simbad database, operated at CDS, Strasbourg, France.

A. THE PRINCIPAL AXES OF THE PROJECTED ELLIPSE

A prolate spheroidal shell centered on the origin, with its major axis aligned with the z axis, is described by the equation:

$$\frac{x^2}{b^2} + \frac{y^2}{b^2} + \frac{z^2}{a^2} = 1$$

with $b < a$. The eccentricity of the ellipse, e , is defined by $b^2 = a^2(1 - e^2)$.

The observer's line of sight, taken to be in the xz plane, makes an angle i with the z axis (polar axis). This observer sees a projected ellipse with semimajor axis $a_\perp$ and semiminor axis $b_\perp$. The prolate symmetry around the z axis gives the result:

$$b_\perp = b = a\sqrt{1 - e^2}.$$

The semimajor projected axis $a_\perp$ can be found using the geometry shown in Figure 1. The intersection of the spheroidal surface and the xz plane is an ellipse described by

$$\frac{x^2}{b^2} + \frac{z^2}{a^2} = 1,$$

or

$$x^2 = a^2(1 - e^2) - z^2(1 - e^2).$$

The line of sight is described generally by

$$z = c + x \cot i.$$

The *tangent* line of sight passes through point A , and for this line c is defined by the condition that the line intersects the ellipse exactly once. Using the equation of the line to substitute for z in the equation of the ellipse, it is seen that

$$x^2[1 + (1 - e^2) \cot^2 i] + x[2c(1 - e^2) \cot i] + (c^2 - a^2)(1 - e^2) = 0$$

This equation, quadratic in x , has a single solution (tangent condition) only when the discriminant, D , is equal to zero:

$$D = 4c^2(1 - e^2)^2 \cot^2 i - 4[1 + (1 - e^2) \cot^2 i](1 - e^2)(c^2 - a^2) = 0.$$

The z -intercept of the line is thus

$$c = a\sqrt{1 + (1 - e^2) \cot^2 i},$$

and the projected semimajor axis is

$$a_{\perp} = c \sin i = a \sin i \sqrt{1 + (1 - e^2) \cot^2 i} = \sqrt{a^2 \sin^2 i + b^2 \cos^2 i} = a \sqrt{1 - e^2 \cos^2 i}.$$

It is easy to see that $a_{\perp} > b_{\perp}$.

The tangent method for finding the projected ellipse was used as long ago as Hubble (1926), although he used it only for oblate spheroids and measured i from the equator rather than the pole of the spheroid.

B. THE MAXIMUM LINE-OF-SIGHT VELOCITY

As before, let the first quadrant of the xz plane contain the observer's line of sight, in a direction defined by the unit vector $\hat{n} = (\sin i, \cos i)$ where i is the angle between the major axis of the spheroid and the observer. By symmetry the maximum projected (line-of-sight) velocity of the ellipsoid will be associated with a point that lies in the xz plane, and it suffices to consider the plane ellipse

$$\frac{x^2}{b^2} + \frac{z^2}{a^2} = 1,$$

or $z^2 = a^2 - x^2(1 - e^2)^{-1}$ where $b^2 = a^2(1 - e^2)$ as before. Let θ be the polar angle defined by $x = z \tan \theta$. (See Figure 2.) Note that

$$2z \frac{dz}{dx} = \frac{d(z^2)}{dx} = -2x(1 - e^2)^{-1},$$

thus

$$\frac{dz}{dx} = -\frac{\tan \theta}{1 - e^2}.$$

A point on the ellipse $\vec{r} = (x, z) = (r \sin \theta, r \cos \theta)$ has velocity $\vec{v} = \vec{r}/t$, where t is the time elapsed since a point explosion. Constant speed (no deceleration) has been assumed. The line-of-sight velocity of gas at this point will be

$$v_{\text{los}} = \vec{v} \cdot \hat{n} = \frac{1}{t}(x \sin i + z \cos i)$$

and the extremum, called v_{max} , occurs for the point $\vec{r}_* = (x_*, z_*)$ such that

$$\frac{dv_{\text{los}}}{dx} = \frac{1}{t} \left(\sin i + \frac{dz}{dx} \cos i \right) = \frac{1}{t} \left(\sin i - \frac{\tan \theta_*}{1 - e^2} \cos i \right) = 0.$$

Thus

$$(1 - e^2) \tan i = \tan \theta_*.$$

Now $\vec{r}_*$ lies on the ellipse, so

$$z_*^2 = a^2 - \frac{x_*^2}{1 - e^2} = a^2 - z_*^2 \frac{\tan^2 \theta_*}{1 - e^2}$$

or after some algebra,

$$z_* = \frac{a}{[1 + (1 - e^2) \tan^2 i]^{1/2}}.$$

After some additional algebra, the desired expression is obtained:

$$v_{\max} = \frac{z_*}{t} (\tan \theta_* \sin i + \cos i) = \frac{a}{t} (1 - e^2 \sin^2 i)^{1/2} = \frac{1}{t} (a^2 \cos^2 i + b^2 \sin^2 i)^{1/2}.$$

Note that in general $\theta_* \neq i$, so that the spot on the projected image of the nova shell where $v_{\max}$ is observed is usually not aligned with the central star.

C. THE ANGLE-AVERAGED APPARENT “RADIUS” OF THE PROJECTED ELLIPSE

Let the projected ellipse be described by

$$\left(\frac{x}{b_{\perp}}\right)^2 + \left(\frac{y}{a_{\perp}}\right)^2 = 1$$

where x, y are now rectangular coordinates in the plane of the sky, and $b_{\perp} < a_{\perp}$. To streamline the notation, the projection subscript ($\perp$) is temporarily suppressed. Using centered polar coordinates (r, θ) such that $y = x \tan \theta$, it can be seen that

$$x^2(a^2 + b^2 \tan^2 \theta) = a^2 b^2,$$

whence

$$\begin{aligned} x^2 &= a^2 b^2 / (a^2 + b^2 \tan^2 \theta) \\ y^2 &= a^2 b^2 \tan^2 \theta / (a^2 + b^2 \tan^2 \theta) \\ r^2 &= x^2 + y^2 = b^2 / [\cos^2 \theta + (b/a)^2 \sin^2 \theta]. \end{aligned}$$

With the projection subscript restored, and with $k' \equiv b_{\perp}/a_{\perp} \leq 1$ and $k^2 \equiv 1 - k'^2$, it can be seen that $r^2(\theta) = b_{\perp}^2 / (1 - k^2 \sin^2 \theta)$. The angle-averaged value of r is thus

$$\bar{r} = \frac{2}{\pi} \int_0^{\pi/2} r(\theta) d\theta = \frac{2b_{\perp}}{\pi} \int_0^{\pi/2} \frac{d\theta}{\sqrt{1 - k^2 \sin^2 \theta}} = \frac{2b_{\perp}}{\pi} K(k) = \frac{2b}{\pi} K(k)$$

where $K(k)$ is the complete elliptic integral of the first kind. By symmetry, the integration is carried out over the first quadrant only.

When $k' \approx 1$, a useful series expansion for $K(k)$ is (e.g. Dwight, 1961)

$$K(k) = \frac{\pi}{2} (1 + m) \left[1 + \frac{1^2}{2^2} m^2 + \frac{1^2 3^2}{2^2 4^2} m^4 + \frac{1^2 3^2 5^2}{2^2 4^2 6^2} m^6 + \dots \right]$$

with $m \equiv (1 - k')/(1 + k') = (a_\perp - b_\perp)/(a_\perp + b_\perp)$. For $k' \geq 0.4$, the series truncated after the m^6 term is accurate to better than three decimal places.

REFERENCES

- Cohen, J.G. 1985, *ApJ*, 292, 90
- Cohen, J.G., & Rosenthal, A.J. 1983, *ApJ*, 268, 689
- Della Valle, M., & Livio, M. 1995, *ApJ*, 452, 704
- Dwight, H. B. 1961, *Tables of Integrals and Other Mathematical Data*, fourth ed. (New Yprk: Macmillan)
- Ford, H. C. & Ciardullo, R. 1988, in *ASP Conf. Ser. 4, The Extragalactic Distance Scale*, ed. S. van den Bergh & C. J. Pritchet (San Francisco: ASP), 128 (FC88)
- Gill, C. D., & O’Brien, T. J. 1998, *MNRAS*, 300, 221
- Gill, C. D., & O’Brien, T. J. 2000, *MNRAS*, in press (astro-ph/0001092)
- Herbig, G. H., & Smak, J. I. 1992, *Acta Astronomica*, 42, 17
- Hjellming, R. M. 1996, in *Cataclysmic Variables and Related Objects*, ed. A. Evans & J. H. Wood (Dordrecht: Kluwer Academic Publishers), 317
- Hjellming, R. M., Wade, C. M., Vandenberg, N. R., & Newell, R. T. 1979, *AJ*, 84, 1619
- Hubble, E. 1926, *ApJ*, 64, 321
- Hutchings, J. B. 1972, *MNRAS*, 158, 177
- Martin, P. G. 1989, in *Classical Novae*, ed. M. F. Bode & A. Evans (Chichester & New York: Wiley), chapter 5.4
- Porter, J. M., O’Brien, T. J., & Bode, M. F. 1998, *MNRAS*, 296, 943
- Seaquist, E. R., Duric, N., Israel, F. P., Spoelstra, T. A. T., Ulich, B. L., & Gregory, P. C. 1980, *AJ*, 85, 283
- Shin, J.-Y., Gehrz, R. D., Jones, T. J., Krautter, J., Heidt, J., & Hjellming, R. M. 1998, *ApJ*, 116, 1966
- Slavin, A. J., O’Brien, T. J., & Dunlop, J. S. 1994, *MNRAS*, 266, L55
- Slavin, A. J., O’Brien, T. J., & Dunlop, J. S. 1995, *MNRAS*, 276, 353
- Soderblom, D. 1976, *PASP*, 88, 517
- Solf, J. 1983, *ApJ*, 273, 647
- Wade, R. A. 1988, 335, 394
- Wade, R. A., & Hubeny, I. 1998, *ApJ*, 509, 350




Fig. 1.— Geometry in the xz plane, used in determining $a_{\perp}$.




Fig. 2.— Geometry for the line-of-sight velocity of a parcel of material in a nova shell.

Table 1. Projection Factors and Distance Bias Factors for Prolate Ellipsoids

i (deg)	k' ($= b_{\perp}/a_{\perp}$)	f_1 ($= v_{\max}/v_0$)	f_2 ($= a_{\perp}/a$)	$\hat{d}_1/d$	$\hat{d}_2/d$	$\hat{d}_3/d$	$\hat{d}_4/d$	$\hat{d}_5/d$
$b/a = f_3 = 1.00$								
0–90	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
$b/a = f_3 = 0.80$								
0.	1.000	1.000	0.800	1.250	1.250	1.250	1.250	1.250
15.	0.982	0.988	0.815	1.212	1.235	1.223	1.223	1.223
30.	0.936	0.954	0.854	1.117	1.192	1.153	1.154	1.154
45.	0.883	0.906	0.906	1.000	1.132	1.062	1.064	1.065
60.	0.839	0.854	0.954	0.896	1.068	0.974	0.978	0.980
75.	0.810	0.815	0.988	0.825	1.019	0.912	0.917	0.919
90.	0.800	0.800	1.000	0.800	1.000	0.889	0.894	0.897
(averaged)	0.858	0.870	0.936	0.937	1.087	1.005	1.009	1.010
$b/a = f_3 = 0.60$								
0.	1.000	1.000	0.600	1.667	1.667	1.667	1.667	1.667
15.	0.945	0.978	0.635	1.541	1.631	1.585	1.585	1.586
30.	0.832	0.917	0.721	1.271	1.528	1.387	1.393	1.396
45.	0.728	0.825	0.825	1.000	1.374	1.158	1.172	1.180
60.	0.655	0.721	0.917	0.787	1.202	0.951	0.972	0.983
75.	0.613	0.635	0.978	0.649	1.058	0.804	0.828	0.841
90.	0.600	0.600	1.000	0.600	1.000	0.750	0.775	0.787
(averaged)	0.695	0.747	0.880	0.886	1.245	1.030	1.048	1.057
$b/a = f_3 = 0.40$								
0.	1.000	1.000	0.400	2.500	2.500	2.500	2.500	2.500
15.	0.860	0.971	0.465	2.089	2.429	2.246	2.252	2.256
30.	0.658	0.889	0.608	1.461	2.222	1.763	1.802	1.822
45.	0.525	0.762	0.762	1.000	1.904	1.311	1.380	1.416
60.	0.450	0.608	0.889	0.684	1.521	0.944	1.020	1.061
75.	0.412	0.465	0.971	0.479	1.163	0.678	0.746	0.783
90.	0.400	0.400	1.000	0.400	1.000	0.571	0.632	0.666
(averaged)	0.506	0.637	0.832	0.861	1.592	1.100	1.161	1.193